

IFT122 Knockout cell line (BEAS-2B)

Catalog Number: KOA23973

Product Information	
Product Name	IFT122 Knockout cell line (BEAS-2B)
specification	1*10 ⁶
Storage and transportation	Shipped on dry ice; Store in liquid nitrogen
Cell morphology	Epithelial-like, adherent
Passage ratio	1:3~1:4
species	Human
Gene	IFT122
Gene ID	55764
Build method	Electroporation/Lentivirus
Mycoplasma testing	negative
Cultivation system	90% DMEM+10% FBS
Price (USD)	Inquiry
Parental Cell Line	BEAS-2B
Quality Control	Genotype: IFT122 Knockout cell line (BEAS-2B)>95% viability before freezing. All cells were tested and found to be free of bacterial, viruses, mycoplasma and other toxins.

Gene Information	
Gene Official Full Name	intraflagellar transport 122 provided by HGNC
Also known as	CED; SPG; CED1; FAP80; WDR10; CFAP80; WDR10p; WDR140
Gene Description	<p>This gene encodes a member of the WD repeat protein family. WD repeats are minimally conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation. This cytoplasmic protein contains seven WD repeats and an AF-2 domain which function by recruiting coregulatory molecules and in transcriptional activation. Mutations in this gene cause cranioectodermal dysplasia-1. A related pseudogene is located on chromosome 3. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2013]</p>

Expression

Broad expression in testis (RPKM 41.7), ovary (RPKM 7.5) and 15 other tissues [See more](#)