

ASAP2 Knockout cell line (BEAS-2B)

Catalog Number: KOA27984

Product Information	
Product Name	ASAP2 Knockout cell line (BEAS-2B)
specification	1*10 ⁶
Storage and transportation	Shipped on dry ice; Store in liquid nitrogen
Cell morphology	Epithelial-like, adherent
Passage ratio	1:3~1:4
species	Human
Gene	ASAP2
Gene ID	8853
Build method	Electroporation/Lentivirus
Mycoplasma testing	negative
Cultivation system	90% DMEM+10% FBS
Price (USD)	Inquiry
Parental Cell Line	BEAS-2B
Quality Control	Genotype: ASAP2 Knockout cell line (BEAS-2B)>95% viability before freezing. All cells were tested and found to be free of bacterial, viruses, mycoplasma and other toxins.

Gene Information	
Gene Official Full Name	ArfGAP with SH3 domain, ankyrin repeat and PH domain 2 provided by HGNC
Also known as	PAP; PAG3; AMAP2; DDEF2; SHAG1; CENTB3; Pap-alpha
Gene Description	This gene encodes a multidomain protein containing an N-terminal alpha-helical region with a coiled-coil motif, followed by a pleckstrin homology (PH) domain, an Arf-GAP domain, an ankyrin homology region, a proline-rich region, and a C-terminal Src homology 3 (SH3) domain. The protein localizes in the Golgi apparatus and at the plasma membrane, where it colocalizes with protein tyrosine kinase 2-beta (PYK2). The encoded protein forms a stable complex with PYK2 in vivo. This interaction appears to be mediated by binding of its SH3 domain to the C-terminal proline-rich domain of PYK2. The encoded protein is tyrosine phosphorylated by activated PYK2. It has catalytic activity for class I and II ArfGAPs in vitro, and can bind the class III Arf ARF6 without immediate GAP activity. The encoded protein is believed to function as an ARF GAP that controls ARF-mediated vesicle budding when recruited to Golgi membranes. In addition, it functions as a substrate

	and downstream target for PYK2 and SRC, a pathway that may be involved in the regulation of vesicular transport. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2008]
Expression	Broad expression in testis (RPKM 24.6), kidney (RPKM 11.9) and 21 other tissues See more