

B4GALT7 Knockout cell line (HCT 116)

Catalog Number: KO01780

Product Information	
Product Name	B4GALT7 Knockout cell line (HCT 116)
specification	1*10^6
Storage and transportation	Dry ice preservation/T25 live cell transportation.
Cell morphology	Epithelioid, adherent cell
Passage ratio	1:2~1:4
species	Human
Gene	B4GALT7
Gene ID	11285
Build method	Electric rotation method / virus method
Mycoplasma testing	Negative
Cultivation system	90%McCOYs 5A+10% FBS
Parental Cell Line	HCT 116
Quality Control	Genotype: B4GALT7 Knockout cell line (HCT 116) >95% viability before freezing. All cells were tested and found to be free of bacterial, viruses,mycoplasma and other toxins.

Gene Information	
Gene Official Full Name	beta-1,4-galactosyltransferase 7provided by HGNC
Also known as	XGPT; EDSP1; XGPT1; EDSSLA; XGALT1; EDSSPD1
Gene Description	This gene is a member of the beta-1,4-galactosyltransferase (beta4GalT) family. Family members encode type II membrane-bound glycoproteins that appear to have exclusive specificity for the donor substrate UDP-galactose. Each beta4GalT member has a distinct function in the biosynthesis of different glycoconjugates and saccharide structures. As type II membrane proteins, they have an N-terminal hydrophobic signal sequence that directs the protein to the Golgi apparatus which then remains uncleaved to function as a transmembrane anchor. The enzyme encoded by this gene attaches the first galactose in the common carbohydrate-protein linkage (GlcA-beta1,3-Gal-beta1,3-Gal-beta1,3-Gal-beta1,4-Xyl-beta1-O-Ser) found in proteoglycans. This enzyme differs from other beta4GalTs because it lacks the conserved Cys residues found in beta4GalT1-beta4GalT6 and it is located in cis-Golgi instead of trans-Golgi. Mutations in this gene have been associated with the progeroid form of Ehlers-Danlos syndrome. [provided by RefSeq, Oct 2009]



Expression

Ubiquitous expression in testis (RPKM 4.7), prostate (RPKM 3.6) and 25 other tissues See more