

HMGA2 Knockout cell line (HeLa)

Catalog Number: KO09870

| Product Information | |
|----------------------------|--|
| Product Name | HMGA2 Knockout cell line (HeLa) |
| specification | 1*10 ⁶ |
| Storage and transportation | Dry ice preservation/T25 live cell transportation. |
| Cell morphology | Epithelioid, adherent cell |
| Passage ratio | 1:3~1:6 |
| species | Human |
| Gene | HMGA2 |
| Gene ID | 8091 |
| Build method | Electric rotation method / virus method |
| Mycoplasma testing | Negative |
| Cultivation system | 90%DMEM+10% FBS |
| Parental Cell Line | HeLa |
| Quality Control | Genotype: HMGA2 Knockout cell line (HeLa) >95% viability before freezing. All cells were tested and found to be free of bacterial, viruses, mycoplasma and other toxins. |

| Gene Information | |
|-------------------------|--|
| Gene Official Full Name | high mobility group AT-hook 2provided by HGNC |
| Also known as | BABL; LIPO; SRS5; HMGIC; HMGI-C; STQTL9 |
| Gene Description | This gene encodes a protein that belongs to the non-histone chromosomal high mobility group (HMG) protein family. HMG proteins function as architectural factors and are essential components of the enhancesome. This protein contains structural DNA-binding domains and may act as a transcriptional regulating factor. Identification of the deletion, amplification, and rearrangement of this gene that are associated with myxoid liposarcoma suggests a role in adipogenesis and mesenchymal differentiation. A gene knock out study of the mouse counterpart demonstrated that this gene is involved in diet-induced obesity. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008] |
| Expression | Low expression observed in reference dataset See more |