

## LAMB1 Knockout cell line (HEK293)

**Catalog Number:** KO10260

Product Information	
Product Name	LAMB1 Knockout cell line (HEK293)
specification	1*10 <sup>6</sup>
Storage and transportation	Dry ice preservation/T25 live cell transportation.
Cell morphology	Epithelioid, adherent cell
Passage ratio	1:3~1:6
species	Human
Gene	LAMB1
Gene ID	3912
Build method	Electric rotation method / virus method
Mycoplasma testing	Negative
Cultivation system	90%DMEM+10% FBS
Parental Cell Line	HEK293
Quality Control	Genotype: LAMB1 Knockout cell line (HEK293) >95% viability before freezing. All cells were tested and found to be free of bacterial, viruses, mycoplasma and other toxins.

Gene Information	
Gene Official Full Name	laminin subunit beta 1 provided by HGNC
Also known as	CLM; LIS5
Gene Description	<p>Laminins, a family of extracellular matrix glycoproteins, are the major noncollagenous constituent of basement membranes. They have been implicated in a wide variety of biological processes including cell adhesion, differentiation, migration, signaling, neurite outgrowth and metastasis.</p> <p>Laminins are composed of 3 non identical chains: laminin alpha, beta and gamma (formerly A, B1, and B2, respectively) and they form a cruciform structure consisting of 3 short arms, each formed by a different chain, and a long arm composed of all 3 chains. Each laminin chain is a multidomain protein encoded by a distinct gene. Several isoforms of each chain have been described. Different alpha, beta and gamma chain isomers combine to give rise to different heterotrimeric laminin isoforms which are designated by Arabic numerals in the order of their discovery, i.e. alpha1beta1gamma1 heterotrimer is laminin 1. The biological functions of the different chains and trimer molecules are largely unknown, but some of the chains have been shown to differ with</p>

	<p>respect to their tissue distribution, presumably reflecting diverse functions in vivo. This gene encodes the beta chain isoform laminin, beta 1. The beta 1 chain has 7 structurally distinct domains which it shares with other beta chain isomers. The C-terminal helical region containing domains I and II are separated by domain alpha, domains III and V contain several EGF-like repeats, and domains IV and VI have a globular conformation. Laminin, beta 1 is expressed in most tissues that produce basement membranes, and is one of the 3 chains constituting laminin 1, the first laminin isolated from Engelbreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chain that is involved in cell attachment, chemotaxis, and binding to the laminin receptor was identified and shown to have the capacity to inhibit metastasis. [provided by RefSeq, Aug 2011]</p>
Expression	Broad expression in placenta (RPKM 75.7), fat (RPKM 65.2) and 23 other tissues See more