

PCSK6 Knockout cell line (A549)

Catalog Number: KO36241

| Product Information | |
|----------------------------|--|
| Product Name | PCSK6 Knockout cell line (A549) |
| specification | 1*10 ⁶ |
| Storage and transportation | Dry ice preservation/T25 live cell transportation. |
| Cell morphology | Epithelioid, adherent cell |
| Passage ratio | 1:3~1:4 |
| species | Human |
| Gene | PCSK6 |
| Gene ID | 5046 |
| Build method | Electric rotation method / virus method |
| Mycoplasma testing | Negative |
| Cultivation system | 90% F12K+10% FBS |
| Parental Cell Line | A549 |
| Quality Control | Genotype: PCSK6 Knockout cell line (A549) >95% viability before freezing. All cells were tested and found to be free of bacterial, viruses, mycoplasma and other toxins. |

| Gene Information | |
|-------------------------|--|
| Gene Official Full Name | proprotein convertase subtilisin/kexin type 6 provided by HGNC |
| Also known as | SPC4; PACE4 |
| Gene Description | <p>This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein undergoes an initial autocatalytic processing event in the ER to generate a heterodimer which exits the ER and sorts to the trans-Golgi network where a second autocatalytic event takes place and the catalytic activity is acquired. The encoded protease is constitutively secreted into the extracellular matrix and expressed in many tissues, including neuroendocrine, liver, gut, and brain. This gene encodes one of the seven basic amino acid-specific members which cleave their substrates at single or paired basic residues. Some of its substrates include transforming growth factor beta related proteins, proalbumin, and von Willebrand factor. This gene is thought to play a role in tumor progression and left-right patterning. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided</p> |

| | |
|------------|--|
| | by RefSeq, Feb 2014] |
| Expression | Biased expression in spleen (RPKM 31.3), liver (RPKM 30.7) and 12 other tissues See more |