

CRYGS Knockout cell line (A549)

Catalog Number: KO38412

Product Information	
Product Name	CRYGS Knockout cell line (A549)
specification	1*10 ⁶
Storage and transportation	Dry ice preservation/T25 live cell transportation.
Cell morphology	Epithelioid, adherent cell
Passage ratio	1:3~1:4
species	Human
Gene	CRYGS
Gene ID	1427
Build method	Electric rotation method / virus method
Mycoplasma testing	Negative
Cultivation system	90% F12K+10% FBS
Parental Cell Line	A549
Quality Control	Genotype: CRYGS Knockout cell line (A549) >95% viability before freezing. All cells were tested and found to be free of bacterial, viruses, mycoplasma and other toxins.

Gene Information	
Gene Official Full Name	crystallin gamma Sprovided by HGNC
Also known as	CRYG8; CTRCT20
Gene Description	<p>Crystallins are separated into two classes: taxon-specific, or enzyme, and ubiquitous. The latter class constitutes the major proteins of vertebrate eye lens and maintains the transparency and refractive index of the lens. Since lens central fiber cells lose their nuclei during development, these crystallins are made and then retained throughout life, making them extremely stable proteins. Mammalian lens crystallins are divided into alpha, beta, and gamma families; beta and gamma crystallins are also considered as a superfamily. Alpha and beta families are further divided into acidic and basic groups. Seven protein regions exist in crystallins: four homologous motifs, a connecting peptide, and N- and C-terminal extensions. Gamma-crystallins are a homogeneous group of highly symmetrical, monomeric proteins typically lacking connecting peptides and terminal extensions. They are differentially regulated after early development. This gene encodes a protein initially considered to be a beta-crystallin but the encoded protein is monomeric and has greater</p>

	sequence similarity to other gamma-crystallins. This gene encodes the most significant gamma-crystallin in adult eye lens tissue. Whether due to aging or mutations in specific genes, gamma-crystallins have been involved in cataract formation. [provided by RefSeq, Jul 2008]
Expression	Ubiquitous expression in spleen (RPKM 1.8), skin (RPKM 1.8) and 25 other tissues See more