

KLRK1 Knockout cell line (AC16)

Catalog Number: KOA10948

Product Information	
Product Name	KLRK1 Knockout cell line (AC16)
specification	1*10 ⁶
Storage and transportation	Shipped on dry ice; Store in liquid nitrogen
Cell morphology	Fibroblast-like, adherent
Passage ratio	1 : 3-1 : 4
species	Human
Gene	KLRK1
Gene ID	22914
Build method	Electroporation/Lentivirus
Mycoplasma testing	negative
Cultivation system	90% DMEM/F12+10% FBS
Price (USD)	Inquiry
Parental Cell Line	AC16
Quality Control	Genotype: KLRK1 Knockout cell line (AC16)>95% viability before freezing. All cells were tested and found to be free of bacterial, viruses, mycoplasma and other toxins.

Gene Information	
Gene Official Full Name	killer cell lectin like receptor K1 provided by HGNC
Also known as	KLR; CD314; NKG2D; NKG2-D; D12S2489E
Gene Description	Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The NKG2 gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed in NK cells. This gene encodes a member of the NKG2 family. The encoded transmembrane protein is characterized by a type II membrane orientation (has an extracellular C terminus) and the presence of a C-type lectin domain. It binds to a diverse family of ligands that include MHC class I chain-related A and B proteins and UL-16 binding proteins, where ligand-receptor interactions can result in the activation of NK and T cells. The surface expression of these ligands is important for the

	recognition of stressed cells by the immune system, and thus this protein and its ligands are therapeutic targets for the treatment of immune diseases and cancers. Read-through transcription exists between this gene and the upstream KLRC4 (killer cell lectin-like receptor subfamily C, member 4) family member in the same cluster. [provided by RefSeq, Dec 2010]
Expression	Broad expression in spleen (RPKM 28.6), lymph node (RPKM 22.6) and 16 other tissues See more