

VEGFA Knockout cell line (AC16)

Catalog Number: KOA13161

Product Information	
Product Name	VEGFA Knockout cell line (AC16)
specification	1*10 ⁶
Storage and transportation	Shipped on dry ice; Store in liquid nitrogen
Cell morphology	Fibroblast-like, adherent
Passage ratio	1 : 3-1 : 4
species	Human
Gene	VEGFA
Gene ID	7422
Build method	Electroporation/Lentivirus
Mycoplasma testing	negative
Cultivation system	90% DMEM/F12+10% FBS
Price (USD)	Inquiry
Parental Cell Line	AC16
Quality Control	Genotype: VEGFA Knockout cell line (AC16)>95% viability before freezing. All cells were tested and found to be free of bacterial, viruses, mycoplasma and other toxins.

Gene Information	
Gene Official Full Name	vascular endothelial growth factor A provided by HGNC
Also known as	VPF; VEGF; MVCD1; L-VEGF
Gene Description	<p>This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome, also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in</p>

	<p>additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site. The levels of VEGF are increased during infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), thus promoting inflammation by facilitating recruitment of inflammatory cells, and by increasing the level of angiopoietin II (Ang II), one of two products of the SARS-CoV-2 binding target, angiotensin-converting enzyme 2 (ACE2). In turn, Ang II facilitates the elevation of VEGF, thus forming a vicious cycle in the release of inflammatory cytokines. [provided by RefSeq, Jun 2020]</p>
Expression	Broad expression in thyroid (RPKM 146.9), prostate (RPKM 69.5) and 21 other tissues See more